

TUMOR SUPPRESSION IN ONCOGENESIS OF ORAL SQUAMOUS CELL CARCINOMA

Elza Ibrahim Auerkari, Agoeng Tjahayani dan Widurini Djaya
Department of Oral Biology, Faculty of Dentistry University of Indonesia

Elza Ibrahim Auerkari, Agoeng Tjahayani dan Widurini Djaya: Tumor Suppression In Oncogenesis Of Oral Squamous Cell Carcinoma. Jurnal Kedokteran Gigi Universitas Indonesia. 2000; 7 (Edisi Khusus): 72-82

Abstract

Oral cancer is relatively uncommon in the industrialized world but accounts for about 5% of all cancer deaths worldwide, and up to 40% of all malignancies in South and South-East Asia. The present paper aims to review the role and significance of tumor suppressor genes in the genetic and molecular pathways to oral squamous cell carcinoma (SCC), which is the most common form of oral cancer and frequently associated with poor prognosis. The high incidence of SCC in betel quid users is due to the severe chemical insult resulting in multiple alterations of oncogenes and tumor suppressing genes. Of the latter, SCC in betel quid users is more often associated with alterations in *p16* and *pRb* (typically at least 60% of the cases) than in *p53* (typically less than 20%). However, in most parts of the world SCC is mostly attributed to smoked tobacco and alcohol, which inflict a synergistic effect when used in combination. The characteristic prevalent alterations are common (50 to 100%) in the tumor suppressor gene *p53*. Potential applications of the genes, corresponding expressed proteins and related other markers are discussed in brief.

Abstrak

Kanker mulut relatif jarang terjadi dinegara industri, namun mencapai sekitar 5% dari seluruh kematian akibat kanker diseluruh dunia, dan mencapai 40% dari seluruh kasus keganasan di Asia selatan dan Asia tenggara. Makalah ini bertujuan untuk membahas peran dan kemaknaan gen supresor tumor pada jalur genetik dan molekuler karsinoma sel skuamosa mulut (SCC), yang merupakan jenis kanker mulut yang paling umum terjadi, dan seringkali dihubungkan dengan prognosis yang buruk. Tingginya insiden SCC pada pemakai pinang, disebabkan oleh iritasi kimia berat yang berakibat terjadinya perubahan-perubahan multipel onkogen dan gen supresor tumor. Dari yang disebut terakhir, SCC pada pengguna pinang lebih sering dihubungkan

perubahan pada p16 dan pRb (secara spesifik paling sedikit 60% dari seluruh kasus) dibandingkan dengan p53 (yang kurang dari 20%). Namun demikian, di sebagian besar belahan dunia, SCC dianggap disebabkan oleh asap tembakau dan alkohol, yang menimbulkan suatu efek sinergistik bila digunakan secara kombinasi. Perubahan-perubahan spesifik yang umum terjadi (50-100%) adalah pada gen supresor p53. Aplikasi penting dari gen-gen, ekspresi protein serta penanda lain yang berhubungan akan dibahas secara singkat.

Introduction

Oral squamous cell carcinoma (SCC) is the most common form of oral cancer, which is the sixth most common malignancy worldwide representing 5.6% of new cancer cases and 5.1% of all cancer deaths.¹ SCC is particularly resistant to therapeutic treatment: about 40-50% of the cases are fatal within 5 years of first diagnosis.² Until recently, there has been little change in the relative survival rates since 1950.³ For further progress, it is clearly important to better understand the mechanisms that lead to oral tumors and control their progression.

Oral cancer is thought to be predominantly initiated and promoted by the chemical insult from oral exposure to certain ecological agents, such as tobacco, ethanol and betel quid.^{4,5} Other contributing though less influential factors appear to include viral infections, poor oral hygiene, consumption of saturated fats and deficiencies in vitamins A and C, and iron.^{6,7,8,9} A rare inherited genetic disorder, Fanconi's anaemia, can also induce oral cancer but probably more important inherited factors are the inter-individual variations in sensitivity to carcinogens.¹⁰ As the patterns of predisposing factors can differ in various parts of the world, so do the incidence and other characteristics of oral cancer. The genetic and molecular pathways in oral cancer can be expected to show corresponding regional differences.

The early premalignant phases of SCC progression result in the development of lesions in the form of leukoplakia (white patches) or erythroplakia (red patches), which can exhibit varying degrees of hyperplasia or dysplasia. Leukoplakia is much more

common but erythroplakia is a more potent precursor of oral squamous cell carcinoma; in fact histopathological examination of the latter type of lesions suggests that about half of these lesions are actually invasive squamous cell carcinomas.¹¹ Of advanced but premalignant dysplastic lesions, 36% on average have been found to transform to a malignant state within 8 years in USA.¹²

There are many oncogenes, or genes which when inappropriately activated, amplified or inactivated can promote some form of cancer: more than 100 oncogenes have been identified.^{13,14} However, cancer is a multistep process where singular gene alterations are necessary but not sufficient for carcinogenesis, and different genetic events are involved in the initiation, promotion, progression and metastasis stages of cancer. Based on epidemiological, molecular and statistical studies it has been suggested that 6 to 10 genetic events are required for oral cancer.¹⁵ In most cases, mechanisms of DNA repair and apoptosis will deal with the deviations from normal cellular function. Cancer is more likely with additional dysfunction, i.e. inappropriate activation or silencing of the tumor suppressing genes (anti-oncogenes) which control some critical events of the cell cycle, including the processes of DNA repair and apoptosis. It is the purpose of the present paper to elucidate the role of tumor suppressor genes in oncogenesis of oral squamous cell carcinomas. Below, a brief review is given on the role of the tumor suppressor genes in the oncogenesis of oral SCC.

Observed Patterns of SCC Oncogenesis and Alteration of Tumor Suppressor Genes

In Europe, USA, Japan, China and Australia the main established etiological agents of oral cancer are smoked tobacco and alcohol (ethanol), which are known to exert a disproportionately stronger effect when used in combination. The synergy is thought to arise sequentially so that carcinogenic and cocarcinogenic chemicals from tobacco induce DNA damage and ethanol promotes its progression by immunosuppressive action and by preventing damage from being repaired. The carcinoma is typically found either endophytically in the trough formed by the lateral border of the tongue, mouth floor and the lingual aspect of the lower alveolus (possibly due to local pooling of carcinogens), or in case of tobacco chewing, exophytically at the site of application, frequently in the buccal sulcus and buccal mucosa. Oral cancer is more common in males than in females, and the incidence rate increases sharply after about 40-50 years of age. In Europe, USA, Japan and Australia about 1-2% of all cancers are oral in origin.¹⁴

In Europe, USA, Japan and China oral carcinomas show a high prevalence of mutations and allelic loss in the important tumor suppressor gene *p53* (Fig 1a), the reported incidence varying generally from about 30% to 95%.¹⁶⁻²⁶ Mutations have been mostly reported from a range of codons from exons 4 to 9, with highest frequencies in hot spots between codons 100 and 300. Particularly common are mutations A→G, G→A, G→T and C→T (Fig 1b). Of these, at least G→T mutations are typical for tobacco derived carcinogens such as benzpyrene.²⁷ The studies reporting highest incidence range (90% or more) of mutations tend to be more recent and to employ direct sequencing of mRNA followed by additional DNA sequencing. The same method also shows higher concordance between primary tumors

and lymph node metastases, and 100% concordance has been reported when using additionally mutation specific oligo ligation assay. This suggests that the *p53* mutations are stable in the progress of cancer.²⁶ Also DNA viruses such as human papilloma virus (HPV) can bind to and potentially inactivate *p53*. As *p53* has multiple important roles (see Table 1) in regulating cell proliferation, DNA repair and apoptosis, its inactivation can lead to cancer in a wide range of tissues including those of the oral cavity.^{13,28} Alterations of *p53* can be already present in pre-malignant lesions, in about 15% of leukoplakias^{21,29} and 30-50% of erythroplakias.³⁰

Frequent inactivation of the tumor suppressing *p16* gene has been reported for primary tumors and metastases of oral SCC and also from about 40% of premalignant lesions.^{31,32} The *p16* protein regulates cell proliferation by binding to cyclin-dependent kinases (CDK) 4 and 6, inhibiting the catalytic activity of the CDK-cyclin D-1 complexes at the G1/S interface by phosphorylation of pRb in the retinoblastoma (Rb) pathway. Allelic loss and mutations of *p16* have been observed in 70-80% of SCC tumors.^{31,33,34} However, in at least 20-30% of the reported cases of SCC, *p16* is not expressed in spite of a normal and intact gene. This is apparently due to DNA methylation at a specific 5' CpG island of the gene. In cultivated SCC cell lines the frequency of altered *p16* can reach 100% but is often observed to be lower in primary tumors. The difference may be partly due to mutations in culture, partly due to techniques used in assessing the altered frequency.³¹ Of the two alternative transcripts (α and β) coded by *p16*, the β transcript gives rise to the p14^{ARF} protein, which by binding to MDM2 leads to stabilization and accumulation of tumor suppressing p53. However, the α transcript which carries the main function of *p16* is much more commonly altered in oral SCC tumors.^{31,35,36}




Fig 1. a) Reported frequency of p53 mutations in SCC tumors in Europe, USA, Japan and China (OECD+China) and in South and South-East Asia (S&SE Asia)^{16-26,39,41-45}; b) frequency of transitions/transversions in human oral SCC tumors (redrawn from the review data of [4]).

A different path of carcinogenesis appears to occur in the well differentiated oral verrucous carcinoma (VC), which shows no metastasis although it is originally described as a subclass of SCC. VC is thought to be initiated by HPV virus²⁵, and the resulting tumor expresses the p16 protein much more strongly (and p53 less) than usual SCC (Fig 2). After emergence of metastasis, the prognosis of SCC is worse but even at this stage there is at least one characteristic tumor suppressing gene, *nm-23*, which is thought to code the NDP kinase involved in transfer of terminal phosphate from a nucleoside triphosphate to NDP.³⁷

The *p16* gene is functionally complementary to *pRb*, which can also be seen as a tumor suppressor gene. In normal cells active pRb prevents cellular transition from G1 to S phase by sequestering the necessary transcription factors. The pRb protein can also bind with the HPV oncoproteins E6 and E7, and high pRb expression together with high levels of p16 is typical in HPV-induced VC.²⁵ In contrast, functional inhibition of p16 but elevated expression of pRb can be observed in usual forms of SCC, particularly in terms of the intensity of the protein expression (Fig 2b).²⁵

Another suggested tumor suppressor gene is *p130*, which is structurally and functionally a relative to *pRb*, and therefore also involved in cell cycle regulation in the G1 phase, with affinity to CDK2. Immunohistochemical activity of p130 is seen in the nuclei and mitochondria.³⁸ Allelic loss in *p130* has been reported in many forms of cancer, and *p130* is apparently particularly affected in poorly differentiated forms of SCC (Fig 3).³⁸

In India, Sri Lanka, Papua New Guinea and parts of Southeast Asia including Indonesia and Taiwan, a particular agent carrying a high risk to oral cancer is chewed tobacco in the form of betel quid. This consists of areca nut (*pinang*) and lime, with or without and tobacco and spices, wrapped into a leaf of the betel vine (*Piper betle*, *sirih*). Exophytical origin of oral cancer is then common, often in the buccal regions. Habitual use of the betel quid is thought to be the main reason for oral cancer in South Asia accounting for 18-40% of all malignancies.^{5,39,40}

Generally less than 20% of the malignant oral tumors have been reported to show mutations in the *p53* gene in South and South East Asia where betel quid is

habitually used (Fig 1a).^{39,41,42,43,44,45} Instead, frequent (35%) mutations of the *H-ras* gene, loss of heterozygosity at the *H-ras* locus (30%), some mutation of the *K-ras* gene (7%), as well as amplification of *N-ras* (28%), *N-myc* (29%), *bcl-2* (23%), *c-myc* (20%) and *K-ras* (17%) oncogenes have been observed.^{39,40,42,46} This multiplicity of genetic alterations appears characteristic for betel quid users, probably reflecting the severity of chemical attack. The *ras* family of oncogenes, often involved early in the formation of chemically induced cancers, result in abnormal G proteins which in normal cells regulate the information exchange across the cell membrane.¹³ The *bcl-2* oncogene expression inhibits both p53-

dependent and p53-independent apoptosis pathways.² The *c-myc* oncogene is involved both in driving the cell cycle and in apoptosis, depending on the level of the growth factor IGF1.⁴⁷ Overexpression of p53 has also been reported but is not well correlated with incidence of mutations.^{45,48} Apparently p53 is inversely expressed with respect to *bcl-2* so that taken together, about 40% of Indian SCC cases may involve dysfunctional apoptosis pathway either via p53 or *bcl-2*.⁴⁰ Perhaps more importantly, frequent loss of both *p16* and *pRb* expression have been reported in SCC tumors and pre-malignant leukoplakias of betel quid users (Fig 2b).⁵




Fig 2 a) Fraction of cells expressing p53, p16 and pRb in common oral SCC tumors, oral verrucous carcinomas (VC), pre-malignant (severe) dysplasias and normal oral tissue, redrawn from data of [25] on Japanese subjects. b) Loss of p16 and pRb expression in SCC tumors, oral premalignant leukoplakias and normal oral tissue from habitual Indian betel/tobacco users (redrawn from data of [5]).




Fig 3. a) Loss of expression of p130 and pRb in oral SCC tumors in all SCC cases and well, moderately and poorly differentiated tumors, and in normal oral tissue (redrawn from data of [38] on Japanese subjects). Note that loss of expression for pRb is much less than that reported for Indian subjects in Fig 2b, with the possible exception of poorly differentiated tumors. b) Frequency of different types of p16 inactivation (mutations in exon or intron, homozygous deletion and methylation) in 14 SCC cell lines (redrawn from data of [31]).

Some of the characteristics of observed oncogenes and tumor suppressor genes associated with oral SCC are summarised in Tables 1 and 2.

Table 1 Chromosomal location and carcinogenetic mechanisms of some recognized oncogenes and anti-oncogenes (tumor suppressor genes) related to oral cancer.^{13,40}

Oncogene	Anti-oncogene	Mechanism ¹⁾	Chromosome
<i>N-ras</i>		IA, PM	1
<i>N-myc</i>		IA	2
<i>K-ras-1</i>		IA, PM	6
<i>c-erb-B-1</i>		IA	7
<i>c-myc</i>		IA, TL	8
	<i>p16</i>	IA, PM, AL, M	9p21-22 (short arm)
<i>H-ras-1, bcl-1, int-2</i>		PM, TL	11
<i>K-ras-2</i>		IA, PM	12
	<i>pRb</i>	I, IA, AL, PM	13
	<i>p130</i>	I, PM, AL	16q12.2-13
	<i>p53</i>	IA, AL, PM	17p13
<i>bcl-2</i>		IA	18q21

1) In carcinogenesis: IA = inappropriate activation/amplification, I = inactivation, PM = point mutation, AL = allelic loss, TL = translocation, M = methylation

Table 2. Features of known tumor suppressor genes associated with oral SCC.^{5,13,25,37}

Gene	Normal function	Notes
<i>p53</i>	Induction of p21 for DNA repair or apoptosis; block of replication of cells with damaged DNA	Altered in many types of tumors including SCC
<i>p16</i>	α -transcript: inactivates cyclin D-dependent kinases 4 and 6 that are needed for phosphorylation of pRb to mediate passage through G1/S	Inactivated (α) in many types of tumors including SCC
<i>p130</i>	Cell cycle regulation, specific for G1 phase, with affinity to CDK2, structural/functional similarity to <i>pRb</i>	Allelic loss in many forms of cancer
<i>pRb</i>	When activated, suppresses cell proliferation by inhibiting G1/S transition via sequestering transcription factors like E2F	Inactivated in many types of tumors including SCC
<i>p23</i>	Encodes NDP kinase involved in transfer of terminal phosphate from a nucleoside triphosphate to NDP	Reduces (in wt- α form) metastatic potential in some cancer types

Other Related Gene/ Protein Markers with Potential Clinical Significance

Apart from those listed above, there is a wide range of cell cycle related proteins that can show important reactions to the status of oral SCC. For example, the antigen protein Ki-67 is a proliferation marker that is active

in all phases of the cell cycle except G₀. Increased Ki-67 expression has been reported as an indicator of proliferation activity (Fig 4a) in pre-malignant and malignant lesions.^{25,49,50} However, the marker efficien-

cy shows considerable scatter and the marker can also fail to reflect the proliferation rate of the tumor, because different antibodies of heterogeneous cell populations recognize different epitopes of the antigen.⁵⁰

The CDK inhibitor p27 has a role in preventing the transition from G1 to S phase.^{25,51} Expression of p 27 generally decreases with increased cell proliferation and is inversely related to Ki-67 expression.²⁵ Significantly elevated apoptotic index has been reported in p27 positive SCC tumors,² and hence p27 may be able to slow down cancer progression. However, generally p27 is less intensively expressed in SCC cells than in normal oral tissue (Fig 4a), and lack of p27 expression may be associated with

poor prognosis.² The corresponding gene *p27* is not sensitive to mutations, but adenovirus E1A and human papillomavirus (HPV) E7 can block the beneficial action of p27.⁵²

The apoptosis regulating protein Bax is a negative regulator of Bcl-2 and is therefore usually inversely distributed in normal tissues with respect to Bcl-2.⁵³ Overexpression of Bax α enhances apoptosis by modulating the p27 protein levels.² Based on experimental evidence on mouse SCC (Fig 4b) it has been suggested that exogenous administering of the *bax* gene in association with chemotherapy may have clinical significance.⁵⁴




Fig 4. a) Fraction of cells expressing the proteins Ki-67 and p27 in oral VC and SCC tumors and in pre-malignant moderate dysplasia (from data of [25]). b) Effect on tumor size of gene gun injection of the *bax* gene with and without chemotherapy (CDDP or cisplatin) initially and after 14 days (4 treatments) in mice inoculated with SCC cells 7 days before therapy (from data of [54]).

Conclusive Remarks

The mechanisms of carcinogenesis of oral SCC are similarly complex as in all cancers, with central role attributed to the accumulating damage inflicted by the etiological agents on the genetic and molecular regulation of the cell cycle. The observed large difference in the incidence of oral SCC between Southern/ Southeastern

Asia and the rest of the world reflects the respective DNA damaging action of the carcinogenic chemicals in chewed betel quid and smoked tobacco. Important part of the damage affects the tumor suppressor genes, of which *p53* and *p16* are particularly sensitive to alteration by mutations, allelic loss and methylation. With the development

efficient analysis techniques, full quantification of the DNA alterations in oral SCC tumors has only recently become possible.

Although many details of the genetic and molecular mechanisms of carcinogenesis and its suppression remain poorly known, the observations made so far hold promise for practical applications both in diagnosis and potential therapy. Further research is needed before new applications can make a significant contribution to the currently dominating therapeutic methods of irradiation, surgery and chemotherapy, which in aggressive combination have recently improved the 5 year survival rates of oral SCC to about 80%.⁵⁵ Positive contribution could be much needed, however, for cases which do not respond well to conventional therapy. In particular, improved survival probability can be expected, if the incidence of complete remission in the therapeutic treatment were to increase.⁵⁵ Current survival rates are lowest and recurrence rates highest in advanced carcinomas with metastasis, which also show the lowest rates of complete remission in the course of therapy. At least to evaluate the potential for carcinogenesis and to discern between primary carcinoma and metastasis, the DNA alterations of the tumor suppressor genes *p53* and *p16* appear promising.^{56,57,58}

The risk of developing primary oral SCC is mainly associated with the habitual use of tobacco, betel quid and alcohol, and the obvious preventive measure is to avoid these carcinogenic agents. The risk of developing second malignancy after treatment for oral SCC tumors has been estimated to be 10-30 times higher than for general population, but recent follow-up studies show that second cancers are likely to appear only in the oral region.^{55,59} This suggests that to prevent secondary tumors, it is necessary to dissuade the patients from continuing to smoking, betel chewing or excessive alcohol consumption.

The new developments have also opened new ways towards cancer chemoprevention, where specific drugs are administered to suppress carcinogenesis and to prevent the development of invasive cancer. Although larger scale confirmation is needed, promising results have been obtained by using 13-*cis*-retinoic acid, which in a limited study resulted in a six-fold decrease in the rate of development of second primary tumors.⁶⁰ Equally important are developments to improve the therapeutic treatments of diagnosed oral SCC. Such promising but as yet experimental methods to enhance the efficiency of chemotherapy include e.g. gene therapy by gene gun injection of the *bax* gene⁵⁴ to the SCC tumors to promote tumor cell apoptosis.

References

1. World Cancer Research Fund, 1997. Food, nutrition and the prevention of cancer: a global perspective. American Institute for Cancer Research, Washington.
2. Fujieda S, Inuzuka M, Tanaka N, Sunaga H, Fan G-K, Ito T *et al.* Expression of p27 is associated with Bax expression and spontaneous apoptosis in oral and oropharyngeal carcinoma. *Int. J. Cancer* 1999; 84: 315-20.
3. Smith CJ. Epidemiology and aetiology. In Langdon JD & Henk JM eds. Malignant tumours of the mouth, jaws and salivary glands. London: Edward Arnold 1995: 1-13.
4. Paterson IC, Eveson JW & Prime SS. Molecular changes in oral cancer may reflect aetiology and ethnic origin. *Oral Oncology* 1996; 32B(3): 150-3.
5. Pande P, Mathur M, Shukla NK & Ralhan R. pRb and p16 protein alterations in human oral tumorigenesis. *Oral Oncology* 1998; 34: 396-403.
6. Stith HF, Rosin MP, Harnby AP, Sankaranarayanan R & Nair MK. Remission of oral leukoplakias and micronuclei in tobacco/betel quid chewers treated with beta-carotene and with beta-carotene plus vitamin A. *Int. J. Cancer* 1988; 421: 195-9.

7. Brachman DG, Graves D & Vokes E. Occurrence of p53 gene deletions and human papilloma virus infection in human head and neck cancer. *Cancer Res* 1992; 52: 4832-6.
8. La Vecchia C, Franceschi S, Levi F, Lucchini F & Negri E. Diet and human carcinoma in Europe. *Oral Oncology* 1993; 29B: 17-22.
9. Fioretti F, Bosetti C, Tavani A, Franceschi S & La Vecchia C. Risk factors for oral and pharyngeal cancer in never smokers. *Oral Oncology* 1999; 35: 375-8.
10. Sato M, Sato T, Izumo T & Amagasa T. Genetic polymorphism of drug-metabolizing enzymes and susceptibility to oral cancer. *Carcinogenesis* 1999; 20(10): 1927-31.
11. Regezi JA, Zarbo RJ, Regev E, Pisanty S, Silverman S & Gazit D. p53 protein expression in sequential biopsies of oral dysplasias and *in situ* carcinomas. *J Oral Pathol Med* 1995; 24: 18-22.
12. Lumerman II, Freedman P & Kerpel S. Oral epithelial dysplasia and the development of invasive squamous cell carcinoma. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* 1995; 79: 321-9.
13. Scully C. Oncogenes, onco-suppressors, carcinogenesis and oral cancer. *Br Dent J* 1992; 173: 53-9.
14. Fox TR & Gonzales AJ. Cell cycle controls as potential targets for the development of chemically induced mouse liver cancer. *CHT Activities* 1996; 16(7): 1-6.
15. Renan MJ. How many mutations are required for tumorigenesis? *Molecular Carcinogenesis* 1993; 7: 139-46.
16. Sakai E & Tsuchida N. Most human squamous cell carcinomas in the oral cavity contain mutated p53 tumor suppressor genes. *Oncogene* 1992; 7: 927-33.
17. Greenblatt MS, Bennett WP, Hollstein M & Harris CC. Mutations in the p53 tumor suppressor gene: clues to cancer etiology and molecular pathogenesis. *Cancer Res* 1994; 53: 4855 - 8.
18. Ahomadegbe JC, Barrois M & Fogel S. High incidence of p53 alterations (mutation, deletion, overexpression) in head and neck primary tumors and metastases; absence of correlation with clinical outcome. Frequent protein overexpression in normal epithelium and in early non-invasive lesions. *Oncogene* 1995;10 : 1217- 27.
19. Kropveld A, van Mansfeld ADM, Nabben N, Hordijk GJ & Slootweg PJ. Discordance of p53 status in matched primary tumors and metastases in head and neck squamous cell carcinoma patients. *Oral Oncology* 1996; 32B: 388-93.
20. Mao EJ, Schwartz SM, Daling JR, Oda D, Tickman L & Beckmann AM. Human papilloma viruses and p53 mutations in normal, premalignant and malignant oral epithelia. *Int J Cancer* 1996; 69: 152-8.
21. Lazarus P, Garewell H, Sciubba J, Zwiebel N, Calcagnotto A, Fair A *et al* . A low incidence of p53 mutations in pre-malignant lesions of the oral cavity from non-tobacco users. *Int. J. Cancer* 1995; 60: 458-63.
22. Mineta H, Borg A, Dictor M, Wahlberg P. & Wennerberg J. Correlation between p53 mutation and cyclin D1 amplification in head and neck squamous cell carcinoma. *Oral Oncology* 1997; 33(1): 42- 6.
23. Shi ST, Yang G-Y, Wang L-D, Xue Z, Feng B, Ding W, Xing EP & Yang CS. 1999. Role of p53 gene expression in human esophageal carcinogenesis: results from immunohistochemical and mutation analyses of carcinomas and nearby non-cancerous lesions. *Carcinogenesis* 1999; 20(4): 591-7.
24. Ibrahim SO, Lillehaug JR, Johannessen AC, Liavaag PG, Nilsen R & Vasstrand EN.. Expression of biomarkers (p53, transforming growth factor alpha, epidermal growth factor receptor, c-erbB-2/neu and the proliferative cell nuclear antigen) in oropharyngeal squamous cell carcinomas. *Oral Oncology* 1999; 35: 302-13.
25. Saito T, Nakajima T & Mogi K. Immunohistochemical analysis of cell cycle-associated proteins p16, pRb, p53, p27 and Ki-67 in oral cancer and precancer with special reference to verrucous carcinomas. *J Oral Pathol Med* 1999; 28: 226-32.
26. Tjebbes GWA, Leppers vd Straat, FGJ, Tilanus MGJ, Hordijk GJ & Slootweg PJ. P53 tumor suppressor gene as a clonal marker in head and neck squamous cell carcinoma: p53 mutations in primary tumor and matched lymph node metastases. *Oral Oncology* 1999; 35: 384-9.

27. Denissenko MF, Pao A, Tang M & Pfeifer G.P. Preferential formation of benzo[α]pyrene adducts at lung cancer mutational hotspots in p53. *Science* 1996; 274: 430-2.
28. Goodger NM, Gannon J, Hunt T & Morgan PR. Cell cycle regulatory proteins - an overview with relevance to oral cancer. *Oral Oncology* 1997; 33(2): 61-73.
29. Boyle JO, Hakim J, Koch W, Van der Riet P, Hruban RH, Roa RA, Corio R., Eby YJ, Ruppert MJ & Sidransky D. The incidence of p53 mutations increases with progression head-and-neck cancer. *Cancer Res* 1993; 53: 4477-80.
30. Qin G-Z, Park JY, Chen S-Y & Lazarus P. A high prevalence of p53 mutations in pre-malignant oral erythroplakia. *Int. J. Cancer* 1999; 80: 345-8.
31. Akanuma D, Uzawa N, Yoshida MA, Negishi A, Amagasa T & Ikeuchi T. Inactivation patterns of the p16 (INK4a) gene in oral squamous cell carcinoma cell lines. *Oral Oncology* 1999; 35: 476-83.
32. Papadimitrakopoulou V, Izzo J, Lippman SM, Lee JS, Fan YH & Clayman G. Frequent inactivation of p16^{INK4a} in oral premalignant lesions. *Oncogene* 1997; 14: 1799-803.
33. Van der Riet P, Nawroz H, Hruban RH, Corio R, Tokino K, Koch W & Sidransky D. Frequent loss of chromosome 9p21-22 early in head and neck cancer progression. *Cancer Research* 1994; 54: 1156-8.
34. Califano J, Van der Riet P, Westra W, Nawroz H, Clayman G, Piantadosi S, Corio R, Lee D, Greenberg B, Koch W & Sidransky D. Genetic progression model for head and neck cancer: implications for field cancerization. *Cancer Research* 1996; 6: 2488-92.
35. Cady II DT, Huang Y, Darnby CJ, Johnson GK & Domann FE. Differential DNA methylation of the p16 INK4/CDKN2A promoter in human oral cancer cells and normal human oral keratinocytes. *Oral Oncology* 1999; 35: 516-22.
36. Hsichischa LI & Reddel RR. p16^{INK4a} and the control of cellular proliferative life span. *Oncogenesis* 1999; 20(6): 921-6.
37. Miyazaki H, Fukuda M, Ishijima Y, Takagi Y, Jimura T, Negishi A, Hirayama R, Iseikawa T. & Kimura N. Overexpression of nm23-H2/NDP Kinase B in a human oral squamous cell carcinoma cell line. in reduced metastasis, differentiated phenotype in the metastatic site, and growth factor-independent proliferative activity in culture. *Clinical Cancer Research* 1999; 5: 4301-7.
38. Tanaka N, Odajima T, Nakano T, Kimijima Y, Yamada S, Ogi K & Kohama G. Immunohistochemical investigation of new suppressor oncogene p130 in oral squamous cell carcinoma. *Oral Oncology* 1999; 35: 321-325.
39. Ranasinghe A, Macgeoch C, Dyer S, Spurr N & Johnson NW. Some oral carcinomas from Sri Lanka betel/tobacco chewers overexpress p53 oncoprotein but lack mutations in exons 5-9. *Anticancer* 1993; 13: 2065-8.
40. Kannan K, Lakshmi Latha PN & Shanmugam G. Expression of bcl-2 oncoprotein in Indian oral squamous cell carcinomas. *Oral Oncology* 1998; 34: 373-7.
41. Chang YS, Lin YJ & Tsai CN. Detection of mutations in the p53 gene in human head and neck carcinomas by single strand conformation polymorphic analysis. *Cancer Lett* 1992; 67: 167-74.
42. Thomas S, Brennan J & Martel G. Mutations in the conserved regions of p53 are infrequent in betel-associated oral cancers from Papua New Guinea. *Cancer Res* 1994; 54: 3588-93.
43. Wong Y-K, Liu T-Y & Chang K-W. p53 alterations in betel quid- and tobacco-associated oral squamous cell carcinomas from Taiwan. *J Oral Pathol Med* 1998; 27: 243-8.
44. Trivedy C, Warnakulasuriya KAAS & Tavassoli M. P53 aberrations in oral submucous fibrosis and oral squamous cell carcinoma detected by immunocytochemistry and PCR-SSCP. *J Oral Pathol Med* 1998; 27: 72-7.
45. Kuo MY-P, Huang J-S, Hsu H-C, Chiang C-P, Kok S-H, Kuo Y-S & Hong C-Y. Infrequent p53 mutations in patients with areca quid chewing-associated oral squamous cell carcinomas in Taiwan. *J Oral Pathol Med* 1999; 28: 221-5.
46. Saranath D, Chang SE & Bhoite LT. High frequency mutations in codons 12 and 61 of H-ras oncogene in chewing tobacco-related human oral carcinoma in India. *Br J Cancer* 1991; 63: 573-8.

47. Hunter T. Oncoprotein networks. *Cell* 1997; 88: 333-46.
48. Kaur J, Srivastava A & Ralhan R. Overexpression of p53 in betel and tobacco related human oral dysplasia and squamous cell carcinoma in India. *Int J Cancer* 1994; 5B: 340-5.
49. Girod SC, Krueger G & Pape HD. 1993. p53 and Ki-67 expression in preneoplastic and neoplastic lesions of the oral mucosa. *Int J Oral Maxillofac Surg* 1993; 22: 285-8.
50. Krecicki T, Jelen M, Zaleska-Krecicka M, Rak J, Szkudlarek T & Jelen-Krzeszewska J. Epidermal growth factor receptor (EGFR), proliferating cell nuclear antigen (PCNA) and Ki-67 antigen in laryngeal epithelial lesions. *Oral Oncology* 1999; 35: 180-6.
51. Sherr CJ & Roberts J. Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev* 1995; 9: 1149-63.
52. Mal A, Poon RY, Howe PH, Toyoshima H, Hunter T & Harter ML. Inactivation of p27Kip1 by the viral E1A oncoprotein in TGFbeta-treated cells. *Nature* 1996; 380: 262-5.
53. Jordan RCK, Catzavelos GC, Barrett AW & Speight PM. Differential expression of bcl-2 and bax in squamous cell carcinomas of the oral cavity. *Oral Oncology* 1996; 32B(6): 394-400.
54. Sugimoto C, Fujieda S, Seki M, Sunaga H, Fan G-K, Tsuzuki H *et al* 1999. Apoptosis-promoting gene (*bax*) transfer potentiates sensitivity of squamous cell carcinoma to cisplatin *in vitro* and *in vivo*. *Int J Cancer* 1999; 82: 860-7.
55. Kirita T, Ohgi K, Shimooka H, Yamanaka Y, Tatebayashi S, Yamamoto K *et al*. Preoperative concurrent chemoradiotherapy plus radical surgery for advanced squamous cell carcinoma of the oral cavity: an analysis of long-term results. *Oral Oncology* 1999; 35: 597-606.
56. Van Oijen, M.G.C.T., Leppers vd Straat F.G.J., Tilanus, M.G.J. & Slootweg, P.J., 2000. The origins of multiple squamous cell carcinomas in the aerodigestive tract. *Cancer* 2000; 88(4): 884-893.
57. Jahnsen S, Karlsson MG. Tumor mapping of regional immunostaining for p21, p53, and mdm2 in locally advanced bladder carcinoma. *Cancer* 2000; 89(3): 619-29.
58. Marin MC, Kaelin Jr WG. P63 and p73: old members of a new family. *Biochim et Biophys Acta* 2000; 1470: M93-M100.
59. Fijuth J, Mazon JJ & Le Pechoux C. Second head and neck cancers following radiation therapy of T1 and T2 cancers of the oral cavity and oropharynx. *Int J Radiation Oncology Biology Physics* 1992; 24: 59-64.
60. Hong WK, Lippman SM & Itri LM. Prevention of second primary tumors with isotretinoin in squamous cell carcinoma of the head and neck. *The New England Journal of Medicine* 1990; 323: 795-801.