

© 2017 Belitung Nursing Journal

This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](#) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited

ORIGINAL RESEARCH

ISSN: 2477-4073

## LAVENDER (*LAVANDULA ANGUSTIFOLIA*) AROMATHERAPY AS AN ALTERNATIVE TREATMENT IN REDUCING PAIN IN PRIMIPAROUS MOTHERS IN THE ACTIVE FIRST STAGE OF LABOR

Hilda Yani Karo Karo\*, Noor Pramono, Sri Wahyuni, Imam Djamaluddin Mashoedi, Leny Latifah

Postgraduate Midwifery Program, Poltekkes Kemenkes Semarang

\*Correspondence:

**Hilda Yani Karo Karo**

Postgraduate Midwifery Program, Poltekkes Kemenkes Semarang

Jl. Tirta Agung, Pedalangan, Banyumanik, Kota Semarang, Jawa Tengah, Indonesia (50268)

E-mail: [hildayanie@rocketmail.com](mailto:hildayanie@rocketmail.com)

### ABSTRACT

**Background:** Labor and childbirth is an extremely painful process. Aromatherapy is considered as one of the nonpharmacological methods to reduce labor pain.

**Objective:** To determine the effect of Lavender (*Lavandula Angustifolia*) aromatherapy on the level of pain in primipara in the first stage of labor.

**Methods:** A quasy-experimental research with pretest and posttest design with control group conducted between October until November 2016. Forty respondents selected using consecutive sampling, which 20 assigned in each group. a Numerical Rating Scale (NRS) pain scale was used. Paired and independent t-test were used for data analyses.

**Results:** The results showed that the p-value of labor pain after intervention was 0.000 (<0.05), which indicated that there was statistically mean difference of labor pain between intervention (6.10) and control group (4.05) in primipara in the first active stage of labor.

**Conclusion:** The women in the lavender aromatherapy group reported lower intensity of labor pain. The intervention study could be practiced in the community health centers for pregnant women in order for them to apply this healing method.

**Key words:** Labor pain, lavender aromatherapy, primipara

### INTRODUCTION

Despite being a natural process, labor and childbirth is an extremely painful process.<sup>1</sup> Many women are fearful and anxious about the pain, which lead to excessive release of hormones such as

catecholamines and steroids that can cause tension of smooth muscle and vasoconstriction of blood vessels.<sup>2</sup> This can decrease uterine contractions, blood flow, and oxygen to the uterus, and the

onset of uterine ischemia that leads to more pain. The long-term pain causes hyperventilation with respiratory frequency 60-70 times per minute, decreases the maternal PaCO<sub>2</sub> level, and increases the pH. If the mother's PaCO<sub>2</sub> levels are low then the fetal PaCO<sub>2</sub> levels will be low too, then resulting in deceleration of the fetal heart rate. In addition, pain causes uncoordinated uterine activity and leads to prolonged labor that can ultimately threaten fetal and maternal life, and leads to increased systole blood pressure resulting in the potential for cardiogenic shock.<sup>3</sup>

Intensity of labor pain in primiparas is often more severe than labor pain in multiparas. It is because multipara mothers experience cervical thinning coinciding with cervical dilatation, while in primipara the cervical thinning process occurs earlier than cervical dilatation.<sup>4</sup> This process causes the intensity of the primiparous contraction to be more severe than multiparous, especially in the first stage of labor. Preliminary survey during October - December 2015 in the working area of the Community Health Center of Delitua showed that there were 42.55% of primigravida had normal duration of the first stage of labor (10-14 hours), 27.65% of primigravida had more than 14 - 20 hours duration, and 10.63% of primigravida had more than 20-24 hours duration of the first stage of labor. Therefore, the intervention to reduce pain that leads to long-term duration of the stage of labor in primipara is needed.

Aromatherapy is one of the interventions considered as an alternative therapy to reduce the pain in primipara. Aromatherapy is a therapy that uses essential oils to help improve or maintain health, encourages, refreshes and awakens body spirit.<sup>5</sup> Lavender (*Lavandula angustifolia*) is one of the essential oil-

producing flowers, which can be used for aromatherapy with the main components of linalool oil (51%) and linalyl acetate (35%).<sup>6</sup> Lavender is also the most popular and safest oil to use, which can increase alpha waves in the brain and help to create a relaxed state and reduce anxiety.<sup>7</sup> Mothers who are undergoing birth, soaking with lavender oil can reduce pain in the perineal area and reduce anxiety.<sup>7</sup>

Research on the effect of aromatherapy using lavender had been studied in Kediri Indonesia, and it is proven effective in reducing pain in mothers during the first stage of labor.<sup>8</sup> However, it is assumed that the aromatherapy might not affect all mothers in Indonesia, due to different taste in using lavender for treatment. Beside, bigger sample size is presented in this study. Therefore, the study aimed to determine the effect of lavender as aromatherapy in reducing pain in mother during the first stage of labor in Deli Serdang, North Sumatera, Indonesia.

## **METHODS**

### *Design*

A quasy experimental research with pretest and posttest design with control group.

### *Setting*

The research was conducted for 2 months between October until November 2016 in the Labor Clinic in the working area of the Community Health Center of Delitua in Deli Serdang Regency.

### *Research subjects*

Forthy respondents selected using consecutive sampling, which 20 assigned in each group. The inclusion criteria included the primigravida in the first active phase of labor (cervix is dilated from 4-10 cm), the height of fundus uteri

was 32 cm - 38 cm above the symphysis, presentation of the back of the head, and willing to be respondent.

#### *Instruments*

A research instrument to measure pain used a Numerical Rating Scale (NRS) pain scale in the range of 0-10, which indicates that 0 refers to no pain, 1-3 refers to mild pain, 4-6 moderate pain, and 7-10 refers to severe pain.<sup>9</sup> Pain was measured before and after intervention.

#### *Intervention*

Lavender essential oil was used as aromatherapy in 10 ml of each pack. The lavender aromatherapy has been applied in the electric aromatherapy diffuser, which was turned on for 5 minutes in an intervention room. When the room was ready, then the researcher invited the respondents to the room for 30 minutes.

#### *Data analysis*

There was no difference in characteristics of the respondents and labor pain with p-value <0.05. Paired and independent t-test were used for data

analysis to determine the influence of lavender aromatherapy on the level of pain in primigravida. Chi square test was also performed for testing confounding factors.

#### *Ethical consideration*

The study permission was obtained from Department of Health of Deli Serdang and the Community Health Center of Delitua. The study has been approved by the Ethical Research Committee with No. 278/KEPK/Poltekkes-Smg/EC/2016. The researchers have confirmed that each respondent have obtained an appropriate informed consent.

## **RESULTS**

Table 1 shows that the average of age of respondents in the intervention and control group was 23 years old, and the cervix dilation was 6. There were no effects of age and cervix dilation on labor pain as the result of chi square with p 0.855 for age, and p 0.314 for cervix dilation (<0.05).

**Table 1. Age and cervix dilation of the respondents and its influence on labor pain**

Group	N	Mean	SD	Min.	Max.	Median	P-value (Chi Square)
<b>Age (Year)</b>							
Intervention	20	23.55	2.16	20	27	24	0.855
Group	20	23.70	2.92	20	29	24	
<b>Cervix Dilation</b>							
Intervention	20	6.10	1.210	4	8	6	0.314
Group	20	5.65	1.599	4	8	5	

**Table 2. Anxiety level in the primigravida mothers and its influence on labor pain using chi-square test**

Group	N	Mild		Moderate		P-value (Chi Square)
		N	%	N	%	
Intervention	20	8	40	12	60	0.114
Group	20	7	35	13	65	

Table 2 shows that 40% of respondents in the intervention group had mild anxiety and 60% had moderate anxiety, while 35% respondents in the control group had mild anxiety and 65%

of them had moderate anxiety. There was no significant influence of anxiety on labor pain with p-value of chi square result 0.114 (<0.05).

**Table 3. Fatigue in the primigravida mothers and its influence on labor pain using chi-square test**

Group	N	Yes		No		P-value (Chi Square)
		N	%	N	%	
Intervention	20	12	60	8	40	0.527
Group	20	10	50	10	50	

As shown in the Table 3, 60% of respondents in the intervention group and 50% in the control group had fatigue.

There was no significant influence of fatigue on labor pain with p-value of chi square result 0.527(<0.05).

**Table 4. Distribution of Labor Pain in the in the intervention and control group using independent t-test**

Pain	Group	Mean	Mean Difference	t	P-value
Pre-test	Control	6.55	-0.30	-0.88	0.381
	Intervention	6.85			
Post-test	Control	6.10	2.05	5.72	0.000
	Intervention	4.05			

Table 4 shows that the mean of pain during pre-test between intervention and control group had no difference with p-value 0.381 (p <0.05), while p-value of pain after intervention was 0.000 (<0.05),

which indicated that there was statistically mean difference of labor pain between intervention (6.10) and control group (4.05).

**Table 5. Effect of lavender aromatherapy on labor pain in the intervention and control group using paired t-test**

Pain	Mean	SD	t	P-value
Control (Pre-Post)	0.450	0.945	2.13	0.046
Intervention (Pre-Post)	2.800	1.196	10.46	0.000

Table 5 shows that the mean of pain before and after intervention in the control group was 0.450 with t count 2.13 and p-value 0.046 (<0.05), while the mean of pain before and after intervention in the intervention group was 2.800 with t count 10.46 and p-value 0.000 (<0.05), which

indicated that there was a significant effect of aromatherapy in the intervention group and standard treatment in control group on labor pain. However, the higher decrease of pain was in the intervention group compared to the decrease of pain in the control group.

## DISCUSSION

Age factor, cervical dilation, anxiety level, and maternal fatigue are often associated with pain before childbirth.<sup>10,11</sup> Therefore, suitable planning and performing midwifery interventions to control the pain during delivery seem to be necessary. However, the result of the present study showed that these factors have not been associated with labor pain.

The results of this study revealed that there is a significant difference of labor pain before and after lavender aromatherapy. This finding supports the finding of the previous study that shows a significant influence of lavender aromatherapy to reduce maternal pain during the first stage of labor.<sup>12</sup> A semi-experimental clinical trial found that women who were treated with lavender aromatherapy during labor reported a lower intensity of pain than women in a control group.<sup>1</sup> Similar with study of Kaviani revealed that The mean of pain intensity perception in the aroma group was lower than that of the control group at 30 and 60 minutes after the aromatherapy intervention ( $p < 0.001$ ).<sup>13</sup>

However, the result of this study is in contrast with the Burns et al study who reported that aromatherapeutic scent had little effect on the pain realisation of the primiparous women.<sup>14</sup> Similar with Yazdkhast revealed that there was no difference in mean duration of the active phase of labor.<sup>15</sup>

Aromatherapy affects the body in three ways: pharmacological effects of the hormones and enzymes that can cause chemical changes in the body; psychological effects, including relaxation and sedative effects on the body; and physiological effects resulting from the brain's response to inhaling aromas.<sup>13,16</sup> The studies conducted on the issue have shown that lavender aroma might suppress

the activity of the sympathetic nervous system.<sup>13,17</sup> Volatile compounds may enter the bloodstream through the nasal or lung mucosa, or directly diffuse into the olfactory nerve and pass up to the limbic system which can influence the sympathetic nervous system.<sup>18</sup>

The results of this study indicated that the lavender aromatherapy is fit with the primiparous women in the North of Sumatera Indonesia and has proven effective in reducing labor pain.

## CONCLUSION

According to the results of this study, the women in the lavender aromatherapy group reported lower intensity of labor pain. This intervention study could be practiced in the community health centers for pregnant women in order for them to apply this healing method.

## REFERENCES

1. Lakhani SE, Sheaffer H, Tepper D. The Effectiveness of Aromatherapy in Reducing Pain: A Systematic Review and Meta-Analysis. *Pain Research and Treatment*. 12/14 09/01/received 11/07/accepted 2016;2016:8158693.
2. Rukiyah AY. Asuhan Kebidanan II (Persalinan). *Jakarta: x*. 2009;195.
3. Astuti R. Pengaruh aromaterapi terhadap nyeri persalinan kala I di bidan praktik swasta kecamatan Polokarto. *Skripsi. Universitas Jenderal Soedirman*. 2009.
4. Winkjosastro GH. Fisiologi janin dalam Ilmu kebidanan: Jakarta, PT Bina Pustaka Sarwono Prawirohardjo; 2008.
5. Bagharpoosh M, Goodarzi GSM. Effect of progressive muscle relaxation technique on pain relief during labor. *Acta Medica Iranica*. 2006;44(3):187-190.

6. Yanti S, Keb M. Buku Ajar Asuhan Kebidanan Persalinan: Yogyakarta: Pustaka; 2010.
7. Cunningham F, Leveno K, Bloom S, Spong CY, Dashe J. *Williams Obstetrics, 24e*: McGraw-Hill; 2014.
8. Kumalasari EP. Studi Tentang Manfaat Aromaterapi (Aroma Lavender) Terhadap Penurunan Tingkat Nyeri Ibu Pada Persalinan Kala I Fase Aktif di Bidan Praktek Swasta Wilayah Kerja Puskesmas Ngletih Kecamatan Pesantren Kota Kediri. *Laporan penelitian. Sekolah Tinggi Ilmu Kesehatan Surya Mitra Husada*. 2012.
9. Judha M. Teori pengukuran nyeri dan nyeri persalinan. *Jogjakarta, Muha Medika*. 2012.
10. Sinambela M. *Pengaruh Tingkat Kecemasan dan Mekanisme Koping Ibu Primigravida terhadap Lamanya Kala I Persalinan Spontan di Klinik Bersalin Swasta Wilayah Kerja Puskesmas Deli Tua Kabupaten Deli Serdang Tahun 2013*.
11. Sulistyowati DID. *Efektifitas terapi aroma lavender terhadap tingkat nyeri dan kecemasan persalinan primipara kala I di rumah sakit dan Klinik Bersalin Purwokerto*, Universitas Indonesia. Fakultas Ilmu Keperawatan; 2009.
12. Karlina SD, Reksohusodo S, Widayati A. Pengaruh Pemberian Aromaterapi Lavender secara Inhalasi terhadap Penurunan Intensitas Nyeri Persalinan Fisiologis pada Primipara Inpartu Kala Satu Fase Aktif di BPM "Fetty Fathiyah" Kota Mataram. *Majalah Kesehatan FKUB*. 2016;2(2):108-119.
13. Kaviani M, Azima S, Alavi N, Tabaei MH. The effect of lavender aromatherapy on pain perception and intrapartum outcome in primiparous women. *British Journal of Midwifery*. 2014;22(2).
14. Burns E, Zobbi V, Panzeri D, Oskrochi R, Regalia A. Aromatherapy in childbirth: a pilot randomised controlled trial. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2007;114(7):838-844.
15. Yazdkhasti M, Pirak A. The effect of aromatherapy with lavender essence on severity of labor pain and duration of labor in primiparous women. *Complement Ther Clin Pract*. Nov 2016;25:81-86.
16. Trevelyan JE, Booth B. *Complementary medicine for nurses, midwives and health visitors*: Macmillan; 1994.
17. Heuberger E, Redhammer S, Buchbauer G. Transdermal absorption of (-)-linalool induces autonomic deactivation but has no impact on ratings of well-being in humans. *Neuropsychopharmacology*. 2004;29(10):1925.
18. Toda M, Morimoto K. Effect of lavender aroma on salivary endocrinological stress markers. *Archives of oral biology*. 2008;53(10):964-968.

**Cite this article as:** Karo HYK, Pramono N, Wahyuni S, Mashoedi ID, Latifah L. Lavender (*Lavandula Angustifolia*) Aromatherapy as an Alternative Treatment in Reducing Pain in Primiparous Mothers in the Active First Stage of Labor. *Belitung Nursing Journal* 2017;3(4):420-425