

Hasil Uji Statistik Menggunakan SPSS

Jenis Kelamin

Crosstabulation

Count

		Diagnosa		Total
		KD	DTK	
JK	Perempuan	9	16	25
	Laki-laki	22	15	37
Total		31	31	62

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	3.284 ^a	1	.070		
Continuity Correction ^b	2.413	1	.120		
Likelihood Ratio	3.319	1	.068		
Fisher's Exact Test				.120	.060
Linear-by-Linear Association	3.231	1	.072		
N of Valid Cases	62				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 12.50.

b. Computed only for a 2x2 table

Usia

Crosstabulation

Count

		DIAG		Total
		KD	DTK	
Usia	0-12 bulan	12	6	18
	13-24 bulan	9	9	18
	25-36 bulan	7	7	14
	37-48 bulan	1	7	8
	48-60 bulan	2	2	4
Total		31	31	62

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	6.500 ^a	4	.165
Likelihood Ratio	7.101	4	.131
Linear-by-Linear Association	3.514	1	.061
N of Valid Cases	62		

a. 4 cells (40.0%) have expected count less than 5. The minimum expected count is 2.00.

Status Gizi Crosstab

Crosstabulation

Count

		DIAG		Total
		KD	DTK	
Gizi	Gizi Kurang	3	4	7
	Gizi Baik	28	27	55
Total		31	31	62

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.161 ^a	1	.688		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.162	1	.688		
Fisher's Exact Test				1.000	.500
Linear-by-Linear Association	.158	1	.691		
N of Valid Cases	62				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 3.50.

b. Computed only for a 2x2 table

Descriptives

		Diagnosa	Statistic	Std. Error	
Hemoglobin	KD	Mean	11.0452	.14575	
		95% Confidence Interval for Mean	Lower Bound	10.7475	
			Upper Bound	11.3428	
		5% Trimmed Mean		11.0649	
		Median		11.3000	
		Variance		.659	
		Std. Deviation		.81152	
		Minimum		9.20	
		Maximum		12.50	
		Range		3.30	
		Interquartile Range		1.30	
		Skewness		-.267	.421
		Kurtosis		-.370	.821
Demam Tanpa Kejang	g	Mean	11.6000	.18883	
		95% Confidence Interval for Mean	Lower Bound	11.2144	
			Upper Bound	11.9856	
		5% Trimmed Mean		11.6190	
		Median		11.8000	

Variance	1.105	
Std. Deviation	1.05135	
Minimum	9.50	
Maximum	13.50	
Range	4.00	
Interquartile Range	1.30	
Skewness	-.585	.421
Kurtosis	-.344	.821

Tests of Normality

	Diagnosa	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Hemo globin	KD	.139	31	.130	.976	31	.682
	Demam Tanpa Kejang	.124	31	.200*	.942	31	.097

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Hemoglobin	Based on Mean	1.454	1	60	.233
	Based on Median	.896	1	60	.348
	Based on Median and with adjusted df	.896	1	55.840	.348
	Based on trimmed mean	1.393	1	60	.242

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Hb	Equal variances assumed	1.454	.233	-2.326	60	.023	-.55484	.23854	-1.03198	-.07769
	Equal variances not assumed			-2.326	56.383	.024	-.55484	.23854	-1.03261	-.07706

Group Statistics

		Diagnosa	N	Mean	Std. Deviation	Std. Error Mean
Hemo globin	KD		31	11.0452	.81152	.14575
	Demam					
	Tanpa Kejang		31	11.6000	1.05135	.18883

Pearson Test

		Hemoglobin	DIAG
Hemoglobin	Pearson Correlation	1	.288*
	Sig. (2-tailed)		.023
	N	62	62
Diagnosa	Pearson Correlation	.288*	1
	Sig. (2-tailed)	.023	
	N	62	62

*. Correlation is significant at the 0.05 level (2-tailed).